arXiv:1301.7057v1 [math.CA] 28 Jan 2013

SOME INTEGRAL INEQUALITIES FOR α-, m-, (α, m)-LOGARITHMICALLY CONVEX FUNCTIONS

MEVLÜT TUNÇ$^{\square}$ AND EBRU YÜKSEL$^{\triangledown}$

Abstract. In this paper, we establish some new Hadamard type inequalities using elementary well known inequalities for functions whose inequalities absolute values are α-, m-, (α, m)-logarithmically convex.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex mapping defined on the interval I of real numbers and $a, b \in I$, with $a < b$. The following double inequalities:

$$f\left(\frac{a+b}{2}\right) \leq \int_a^b f(x)\,dx \leq \frac{f(a)+f(b)}{2}$$

hold. This double inequality is known in the literature as the Hermite-Hadamard inequality for convex functions (see [1]-[8]).

In this section, we will present definitions and some results used in this paper.

Definition 1. *Let I be an interval in $\mathbb{R}$. Then $f : I \to \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$ is said to be convex if*

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y). \tag{1.1}$$

for all $x, y \in I$ and $t \in [0,1]$.

In [1], the concepts of α-, m- and (α, m)-logarithmically convex functions were introduced as follows.

Definition 2. [1] *A function $f : [0,b] \to (0,\infty)$ is said to be m-logarithmically convex if the inequality*

$$f(tx + m(1-t)y) \leq [f(x)]^t [f(y)]^{m(1-t)} \tag{1.2}$$

holds for all $x, y \in [0,b]$, $m \in (0,1]$, and $t \in [0,1]$.

Obviously, if putting $m = 1$ in Definition 2, then f is just the ordinary logarithmically convex on $[0,b]$.

Definition 3. [8] *A function $f : [0,b] \to (0,\infty)$ is said to be α-logarithmically convex if*

$$f(tx + (1-t)y) \leq [f(x)]^{t^\alpha} [f(y)]^{(1-t^\alpha)} \tag{1.3}$$

Date: January 10, 2013.

2000 *Mathematics Subject Classification.* 26A15, 26A51, 26D10.

Key words and phrases. α-, m-, (α, m)-logarithmically convex, Hadamard's inequality, Hölder's inequality, power mean inequality, Cauchy's inequality.

$^{\square}$Corresponding Author.

This paper is in final form and no version of it will be submitted for publication elsewhere.

holds for all $x, y \in [0, b]$, $\alpha \in (0, 1]$ *and* $t \in [0, 1]$.

Clearly, when taking $\alpha = 1$ in Definition 3, then f becomes the ordinary logarithmically convex on $[0, b]$.

Definition 4. [1] *A function* $f : [0, b] \to (0, \infty)$ *is said to be* (α, m)*-logarithmically convex if*

$$f\left(tx + m\left(1 - t\right)y\right) \leq \left[f\left(x\right)\right]^{t^{\alpha}}\left[f\left(y\right)\right]^{m\left(1 - t^{\alpha}\right)} \tag{1.4}$$

holds for all $x, y \in [0, b]$, $(\alpha, m) \in (0, 1] \times (0, 1]$, *and* $t \in [0, 1]$.

Clearly, when taking $\alpha = 1$ in Definition 4, then f becomes the standard m-logarithmically convex function on $[0, b]$, and, when taking $m = 1$ in Definition 4, then f becomes the α-logarithmically convex function on $[0, b]$.

2. NEW HADAMARD-TYPE INEQUALITIES

In order to prove our main theorems, we need the following lemma [2].

Lemma 1. [2] *Let* $f : I \subset \mathbb{R} \to \mathbb{R}$ *be a differentiable mapping on* I°, $a, b \in I^{\circ}$ *with* $a < b$. *If* $f' \in L\left[a, b\right]$, *then the following equality holds:*

$$\begin{aligned} &\frac{f\left(a\right) + f\left(b\right)}{2} - \frac{1}{b - a}\int_{a}^{b} f\left(x\right)dx \\ = &\frac{b - a}{2}\int_{0}^{1}\int_{0}^{1}\left[f'\left(ta + \left(1 - t\right)b\right) - f'\left(sa + \left(1 - s\right)b\right)\right]\left(s - t\right)dtds. \end{aligned} \tag{2.1}$$

A simple proof of this equality can be also done integrating by parts in the right hand side (see [2]).

The next theorems gives a new result of the upper Hermite-Hadamard inequality for α-, m-, (α, m)-logarithmically convex functions.

Theorem 1. *Let* $I \supset [0, \infty)$ *be an open interval and let* $f : I \to (0, \infty)$ *be a differentiable function on* I *such that* $f' \in L\left(a, b\right)$ *for* $0 \leq a < b < \infty$. *If* $\left|f'\left(x\right)\right|$ *is* (α, m)*-logarithmically convex on* $\left[0, \frac{b}{m}\right]$ *for* $(\alpha, m) \in (0, 1]^2$, *then*

$$\begin{aligned} &\left|\frac{f\left(a\right) + f\left(b\right)}{2} - \frac{1}{b - a}\int_{a}^{b} f\left(x\right)dx\right| \\ \leq &\begin{cases} \frac{(b-a)}{3}\left|f'\left(\frac{b}{m}\right)\right|^{m}, & \eta = 1 \\ \frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^{m}\frac{-\alpha^2\ln^2\eta - 2\alpha\ln\eta + 2\eta^{\alpha} - 2}{\alpha^3\ln^3\eta}, & \eta < 1 \end{cases} \end{aligned} \tag{2.2}$$

where $\eta = \left|f'\left(a\right)\right| / \left|f'\left(\frac{b}{m}\right)\right|^{m}$.

Proof. By Lemma 1 and since $|f'|$ is an (α, m)-logarithmically convex on $\left[0, \frac{b}{m}\right]$, then we have

$$
\begin{aligned}
& \left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\
\leq\ & \frac{b-a}{2}\int_0^1\int_0^1 \left|\left(f'(ta+(1-t)b)\right)-\left(f'(sa+(1-s)b)\right)\right| |s-t|\,dtds \\
\leq\ & \frac{b-a}{2}\left[\int_0^1\int_0^1 |s-t|\,|f'(a)|^{t^\alpha}\left|f'\left(\frac{b}{m}\right)\right|^{m(1-t^\alpha)} dtds\right] \\
& +\frac{b-a}{2}\left[\int_0^1\int_0^1 |s-t|\,|f'(a)|^{s^\alpha}\left|f'\left(\frac{b}{m}\right)\right|^{m(1-s^\alpha)} dtds\right]
\end{aligned}
$$

If $0<k\leq 1$, $0<m,n\leq 1$

$$k^{m^n}\leq k^{mn}. \tag{2.3}$$

When $\eta=1$, by (2.3), we get

$$
\begin{aligned}
& \left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\
\leq\ & \frac{b-a}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left[\int_0^1\int_0^1 |s-t|\,dtds+\int_0^1\int_0^1 |s-t|\,dtds\right] \\
=\ & \frac{b-a}{3}\left|f'\left(\frac{b}{m}\right)\right|^m
\end{aligned}
$$

When $\eta<1$, by (2.3), we get

$$
\begin{aligned}
& \left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\
\leq\ & \frac{b-a}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left[\int_0^1\int_0^1 |s-t|\,\eta^{\alpha t}dtds+\int_0^1\int_0^1 |s-t|\,\eta^{\alpha s}dtds\right] \\
=\ & \frac{b-a}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left[\frac{-\alpha^2\ln^2\eta-2\alpha\ln\eta+4\eta^\alpha+\alpha^2\eta^\alpha\ln^2\eta-2\alpha\eta^\alpha\ln\eta-4}{2\alpha^3\ln^3\eta}\right. \\
& \left.+\frac{-\alpha\ln\eta+2\eta^\alpha-\alpha\eta^\alpha\ln\eta-2}{2\alpha^2\ln^2\eta}\right]
\end{aligned}
$$

which completes the proof. □

Corollary 1. *Let $I\supset[0,\infty)$ be an open interval and let $f: I\to(0,\infty)$ be a differentiable function on I such that $f'\in L(a,b)$ for $0\leq a<b<\infty$. If $|f'(x)|$ is m-logarithmically convex on $\left[0,\frac{b}{m}\right]$ for $m\in(0,1]$, then*

$$
\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\leq\begin{cases}\frac{(b-a)}{3}\left|f'\left(\frac{b}{m}\right)\right|^m, & \eta=1\\ \frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\frac{-\ln^2\eta-2\ln\eta+2\eta-2}{\ln^3\eta}, & \eta<1\end{cases}
$$

where η is same as Theorem 1.

Corollary 2. *Let $I\supset[0,\infty)$ be an open interval and let $f:\ I\rightarrow(0,\infty)$ be a differentiable function on I such that $f'\in L(a,b)$ for $0\leq a<b<\infty$. If $|f'(x)|$is α-logarithmically convex on $[0,b]$ for $\alpha\in(0,1]$, then*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\leq\left\{\begin{array}{ll}\frac{(b-a)}{3}\left|f'(b)\right|, & \eta=1\\ \frac{(b-a)}{2}\left|f'(b)\right|\frac{4\eta^{\alpha}-4\alpha\ln\eta-2\alpha^2\ln^2\eta-4}{2\alpha^3\ln^3\eta}, & \eta<1\end{array}\right.$$

where $\eta=|f'(a)|/|f'(b)|$.

Theorem 2. *Let $I\supset[0,\infty)$ be an open interval and let $f:\ I\rightarrow(0,\infty)$ be a differentiable function on I such that $f'\in L(a,b)$ for $0\leq a<b<\infty$. If $|f'(x)|^q$ is an (α,m)-logarithmically convex on $\left[0,\frac{b}{m}\right]$ for $(\alpha,m)\in(0,1]^2$ and $p,q>1$ with $\frac{1}{p}+\frac{1}{q}=1$, then*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \tag{2.4}$$
$$\leq\left\{\begin{array}{ll}(b-a)\left|f'\left(\frac{b}{m}\right)\right|^m\left(\frac{2}{(p+1)(p+2)}\right)^{\frac{1}{p}}, & \eta=1\\ (b-a)\left|f'\left(\frac{b}{m}\right)\right|^m\left(\frac{2}{(p+1)(p+2)}\right)^{\frac{1}{p}}\times\left(\frac{\eta(\alpha q,\alpha q)-1}{\ln\eta(\alpha q,\alpha q)}\right)^{\frac{1}{q}}, & \eta<1\end{array}\right.$$

where $\eta(\alpha,\alpha)$ is same as Theorem 1

Proof. Since $|f'|^q$ is an (α,m)-logarithmically convex on $\left[0,\frac{b}{m}\right]$, from Lemma 1 and the well known Hölder inequality, we have

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\\ \leq\ &\frac{b-a}{2}\int_0^1\int_0^1\left|\left(f'(ta+(1-t)b)\right)-\left(f'(sa+(1-s)b)\right)\right||s-t|\,dtds\\ \leq\ &\frac{b-a}{2}\int_0^1\int_0^1|s-t|\left|f'(a)\right|^{t^{\alpha}}\left|f'\left(\frac{b}{m}\right)\right|^{m(1-t^{\alpha})}dtds\\ &+\frac{b-a}{2}\int_0^1\int_0^1|s-t|\left|f'(a)\right|^{s^{\alpha}}\left|f'\left(\frac{b}{m}\right)\right|^{m(1-s^{\alpha})}dtds\\ \leq\ &\frac{b-a}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1|s-t|^p\,dtds\right)^{\frac{1}{p}}\\ &\times\left[\left(\int_0^1\int_0^1\eta^{qt^{\alpha}}dtds\right)^{\frac{1}{q}}+\left(\int_0^1\int_0^1\eta^{qs^{\alpha}}dtds\right)^{\frac{1}{q}}\right]\end{aligned} \tag{2.5}$$

If $\eta = 1$, by (2.3), we obtain

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx \right| \\ \leq & \quad (b-a)\left| f'\left(\frac{b}{m}\right)\right|^m \left(\int_0^1\int_0^1 |s-t|^p \,dtds\right)^{\frac{1}{p}} \\ = & \quad (b-a)\left| f'\left(\frac{b}{m}\right)\right|^m \left(\frac{2}{(p+1)(p+2)}\right)^{\frac{1}{p}} \end{aligned}$$

If $\eta < 1$, by (2.3), we obtain

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx \right| \\ \leq & \quad \frac{b-a}{2}\left| f'\left(\frac{b}{m}\right)\right|^m \left(\int_0^1\int_0^1 |s-t|^p \,dtds\right)^{\frac{1}{p}} \\ & \times \left[\left(\int_0^1\int_0^1 \eta^{qt^\alpha} dtds\right)^{\frac{1}{q}} + \left(\int_0^1\int_0^1 \eta^{qs^\alpha} dtds\right)^{\frac{1}{q}}\right] \\ = & \quad (b-a)\left| f'\left(\frac{b}{m}\right)\right|^m \left(\frac{2}{(p+1)(p+2)}\right)^{\frac{1}{p}} \times \left(\frac{\eta(\alpha q, \alpha q)-1}{\ln \eta(\alpha q,\alpha q)}\right)^{\frac{1}{q}} \end{aligned} \tag{2.6}$$

which completes the proof. □

Corollary 3. *Let $I \supset [0,\infty)$ be an open interval and let $f : I \to (0,\infty)$ be a differentiable function on I such that $f' \in L(a,b)$ for $0 \leq a < b < \infty$. If $|f'(x)|^q$ is an m-logarithmically convex on $\left[0, \frac{b}{m}\right]$ for $m \in (0,1]$ and $p = q = 2$, then*

$$\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx \right| \leq \begin{cases} (b-a)\left|f'\left(\frac{b}{m}\right)\right|^m \sqrt{\frac{1}{6}}, & \eta = 1 \\ (b-a)\left|f'\left(\frac{b}{m}\right)\right|^m \sqrt{\frac{1}{6}}\left(\frac{\eta(2,2)-1}{\ln \eta(2,2)}\right)^{\frac{1}{2}}, & \eta < 1 \end{cases}$$

Corollary 4. *Let $I \supset [0,\infty)$ be an open interval and let $f : I \to (0,\infty)$ be a differentiable function on I such that $f' \in L(a,b)$ for $0 \leq a < b < \infty$. If $|f'(x)|$ is α-logarithmically convex on $[0,b]$ for $\alpha \in (0,1]$, then*

$$\begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx \right| \\ \leq & \begin{cases} (b-a)|f'(b)|\left(\frac{2}{(p+1)(p+2)}\right)^{\frac{1}{p}}, & \eta = 1 \\ (b-a)|f'(b)|\left(\frac{2}{(p+1)(p+2)}\right)^{\frac{1}{p}}\left(\frac{\eta(\alpha q,\alpha q)-1}{\ln \eta(\alpha q,\alpha q)}\right)^{\frac{1}{q}}, & \eta < 1 \end{cases} \end{aligned}$$

where $\eta = |f'(a)| / |f'(b)|$.

Theorem 3. *Let $I \supset [0,\infty)$ be an open interval and let $f : I \to (0,\infty)$ be a differentiable function on I such that $f' \in L(a,b)$ for $0 \leq a < b < \infty$. If $|f'(x)|^q$ is*

(α, m)-logarithmically convex on $\left[0, \frac{b}{m}\right]$ for $(\alpha, m) \in (0,1]^2$, and then

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \leq \begin{cases} \frac{b-a}{3}\left|f'\left(\frac{b}{m}\right)\right|^m, & \eta=1 \\ \frac{b-a}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m\left\{\left[\frac{2\varphi-2}{[\ln\varphi]^3}-\frac{\varphi+1}{[\ln\varphi]^2}-\frac{1-\varphi}{2\ln\varphi}\right]^{\frac{1}{q}}+\left[\frac{\varphi-1}{[\ln\varphi]^2}-\frac{\varphi+1}{2\ln\varphi}\right]^{\frac{1}{q}}\right\}, & \eta<1 \end{cases} \tag{2.7}$$

where $\eta(\alpha,\alpha)$ is same as Theorem 1, and we take $\eta(\alpha q, \alpha q)=\varphi$.

Proof. Since $|f'|^q$ is an (α, m)-logarithmically convex on $\left[0, \frac{b}{m}\right]$, for $q \geq 1$, from Lemma 1 and the well known power mean integral inequality, we get

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\ \leq\ & \frac{b-a}{2}\int_0^1\int_0^1 \left|\left(f'(ta+(1-t)b)\right)-\left(f'(sa+(1-s)b)\right)\right||s-t|\,dtds \\ \leq\ & \frac{b-a}{2}\left(\int_0^1\int_0^1|s-t|\,dtds\right)^{1-\frac{1}{q}}\left(\int_0^1\int_0^1|s-t|\left|f'(ta+(1-t)b)\right|^q dtds\right)^{\frac{1}{q}} \\ & +\frac{b-a}{2}\left(\int_0^1\int_0^1|s-t|\,dtds\right)^{1-\frac{1}{q}}\left(\int_0^1\int_0^1|s-t|\left|f'(sa+(1-s)b)\right|^q dtds\right)^{\frac{1}{q}} \\ \leq\ & \frac{b-a}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1|s-t|\,dtds\right)^{1-\frac{1}{q}}\left(\int_0^1\int_0^1|s-t|\,\eta^{qt^\alpha}dtds\right)^{\frac{1}{q}} \\ & +\frac{b-a}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1|s-t|\,dtds\right)^{1-\frac{1}{q}}\left(\int_0^1\int_0^1|s-t|\,\eta^{qs^\alpha}dtds\right)^{\frac{1}{q}} \end{aligned}$$

When $\eta=1$, by (2.3), we obtain

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\ \leq\ & \frac{b-a}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1|s-t|\,dtds\right)^{\frac{1}{q}} \\ & +\frac{b-a}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1|s-t|\,dtds\right)^{\frac{1}{q}} \\ =\ & \frac{b-a}{3}\left|f'\left(\frac{b}{m}\right)\right|^m \end{aligned}$$

When $\eta < 1$, by (2.3), we obtain

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\
\leq\ & \frac{b-a}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1 |s-t|\,\eta^{\alpha q t}dtds\right)^{\frac{1}{q}} \\
&+\frac{b-a}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m\left(\int_0^1\int_0^1 |s-t|\,\eta^{\alpha q s}dtds\right)^{\frac{1}{q}} \\
=\ & \frac{b-a}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m \\
&\times\left\{\left[\frac{2\eta(\alpha q,\alpha q)-2}{[\ln(\eta(\alpha q,\alpha q))]^3}-\frac{\eta(\alpha q,\alpha q)+1}{[\ln(\eta(\alpha q,\alpha q))]^2}-\frac{1-\eta(\alpha q,\alpha q)}{2\ln(\eta(\alpha q,\alpha q))}\right]^{\frac{1}{q}}\right. \\
&\left.+\left[\frac{\eta(\alpha q,\alpha q)-1}{[\ln(\eta(\alpha q,\alpha q))]^2}-\frac{\eta(\alpha q,\alpha q)+1}{2\ln(\eta(\alpha q,\alpha q))}\right]^{\frac{1}{q}}\right\}
\end{aligned}$$

which completes the proof. □

Corollary 5. *Let $I \supset [0,\infty)$ be an open interval and let $f : I \to (0,\infty)$ be a differentiable function on I such that $f' \in L(a,b)$ for $0 \leq a < b < \infty$. If $|f'(x)|^q$ is m-logarithmically convex on $\left[0,\frac{b}{m}\right]$ for $m \in (0,1]$, then*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|$$

$$\leq\begin{cases}\frac{b-a}{3}\left|f'\left(\frac{b}{m}\right)\right|^m, & \eta=1\\ \frac{(b-a)}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}\left|f'\left(\frac{b}{m}\right)\right|^m \\ \quad\times\left\{\left[\frac{2\eta(q,q)-2}{[\ln\eta(q,q)]^3}-\frac{\eta(q,q)+1}{[\ln\eta(q,q)]^2}-\frac{1-\eta(q,q)}{2\ln\eta(q,q)}\right]^{\frac{1}{q}}+\left[\frac{\eta(q,q)-1}{[\ln\eta(q,q)]^2}-\frac{\eta(q,q)+1}{2\ln\eta(q,q)}\right]^{\frac{1}{q}}\right\}, & \eta<1\end{cases}$$

Corollary 6. *Let $I \supset [0,\infty)$ be an open interval and let $f : I \to (0,\infty)$ be a differentiable function on I such that $f' \in L(a,b)$ for $0 \leq a < b < \infty$. If $|f'(x)|$ is α-logarithmically convex on $[0,b]$ for $\alpha \in (0,1]$, then*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|$$

$$\leq\begin{cases}(b-a)\left(\frac{1}{3}\right)|f'(b)|, & \eta=1\\ \frac{(b-a)}{2}\left(\frac{1}{3}\right)^{1-\frac{1}{q}}|f'(b)|\left(\left[\frac{2\eta(\alpha q,\alpha q)-2}{[\ln(\eta(\alpha q,\alpha q))]^3}-\frac{\eta(\alpha q,\alpha q)+1}{[\ln(\eta(\alpha q,\alpha q))]^2}-\frac{1-\eta(\alpha q,\alpha q)}{2\ln(\eta(\alpha q,\alpha q))}\right]^{\frac{1}{q}}\right. \\ \quad\left.+\left[\frac{\eta(\alpha q,\alpha q)-1}{[\ln(\eta(\alpha q,\alpha q))]^2}-\frac{\eta(\alpha q,\alpha q)+1}{2\ln(\eta(\alpha q,\alpha q))}\right]^{\frac{1}{q}}\right), & \eta<1\end{cases}$$

where $\eta = |f'(a)| / |f'(b)|$.

Theorem 4. *Let $f : I \subset \mathbb{R}_+ \to \mathbb{R}_+$ be differentiable on I°, $a, b \in I$, with $a < b$ and $f' \in L([a,b])$. If $|f'|$ is an (α,m)-logarithmically convex $\left[0,\frac{b}{m}\right]$ for $(\alpha,m) \in (0,1]^2$*

and $\mu_1, \mu_2, \tau_1, \tau_2 > 0$ *with* $\mu_1 + \tau_1 = 1$ *and* $\mu_2 + \tau_2 = 1$*, then*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx\right| \tag{2.8}$$

$$\leq \begin{cases} \frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^m \left[\frac{2\mu_1^3}{(2\mu_1+1)(\mu_1+1)} + \frac{2\mu_2^3}{(2\mu_2+1)(\mu_2+1)} + \tau_1 + \tau_2\right], & \eta = 1 \\ \frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^m \left\{\frac{2\mu_1^3}{(2\mu_1+1)(\mu_1+1)} + \frac{2\mu_2^3}{(2\mu_2+1)(\mu_2+1)} \right. & \\ \qquad \left. + \tau_1 \frac{\eta\left(\frac{\alpha}{\tau_1}, \frac{\alpha}{\tau_1}\right) - 1}{\ln \eta\left(\frac{\alpha}{\tau_1}, \frac{\alpha}{\tau_1}\right)} + \tau_2 \frac{\eta\left(\frac{\alpha}{\tau_2}, \frac{\alpha}{\tau_2}\right) - 1}{\ln \eta\left(\frac{\alpha}{\tau_2}, \frac{\alpha}{\tau_2}\right)}\right\}, & \eta < 1 \end{cases}$$

where $\eta(\alpha, \alpha)$ *is same as Theorem 1.*

Proof. Since $|f'|^q$ is an (α, m)-logarithmically convex on $\left[0, \frac{b}{m}\right]$, from Lemma 1, we have

$$\begin{aligned} &\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx\right| \\ \leq\ & \frac{(b-a)}{2}\int_0^1\int_0^1 \left|(f'(ta+(1-t)b)) - (f'(sa+(1-s)b))\right| |s-t|\, dtds \\ \leq\ & \frac{(b-a)}{2}\left[\int_0^1\int_0^1 |s-t| \left|f'(a)\right|^{t^\alpha} \left|f'\left(\frac{b}{m}\right)\right|^{m(1-t^\alpha)} dtds\right] \\ & + \frac{(b-a)}{2}\left[\int_0^1\int_0^1 |s-t| \left|f'(a)\right|^{s^\alpha} \left|f'\left(\frac{b}{m}\right)\right|^{m(1-s^\alpha)} dtds\right] \\ =\ & \frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^m \left[\int_0^1\int_0^1 |s-t|\,\eta^{t^\alpha} dtds + \int_0^1\int_0^1 |s-t|\,\eta^{s^\alpha} dtds\right] \end{aligned} \tag{2.9}$$

for all $t \in [0,1]$. Using the well known inequality $rt \leq \mu r^{\frac{1}{\mu}} + \tau t^{\frac{1}{\tau}}$, on the right side of (2.9), we get

$$\begin{aligned} & \int_0^1\int_0^1 |s-t|\,\eta^{t^\alpha} dtds + \int_0^1\int_0^1 |s-t|\,\eta^{s^\alpha} dtds \\ \leq\ & \mu_1 \int_0^1\int_0^1 |s-t|^{\frac{1}{\mu_1}} dtds + \tau_1 \int_0^1\int_0^1 \eta^{\frac{t^\alpha}{\tau_1}} dtds \\ & + \mu_2 \int_0^1\int_0^1 |s-t|^{\frac{1}{\mu_2}} dtds + \tau_2 \int_0^1\int_0^1 \eta^{\frac{s^\alpha}{\tau_2}} dtds \end{aligned} \tag{2.10}$$

When $\eta = 1$, by (2.3), we get

$$\begin{aligned} & \int_0^1\int_0^1 |s-t|\,\eta^{t^\alpha} dtds + \int_0^1\int_0^1 |s-t|\,\eta^{s^\alpha} dtds \\ \leq\ & \frac{2\mu_1^3}{(2\mu_1+1)(\mu_1+1)} + \frac{2\mu_2^3}{(2\mu_2+1)(\mu_2+1)} + \tau_1 + \tau_2 \end{aligned} \tag{2.11}$$

When $\eta < 1$, by (2.3), we get

$$\begin{aligned}
&\int_0^1\int_0^1 |s-t|\,\eta^{t^{\alpha}}dtds+\int_0^1\int_0^1 |s-t|\,\eta^{s^{\alpha}}dtds \\
\leq\ & \mu_1\int_0^1\int_0^1 |s-t|^{\frac{1}{\mu_1}}dtds+\tau_1\int_0^1\int_0^1 \eta^{\frac{t^{\alpha}}{\tau_1}}dtds+\mu_2\int_0^1\int_0^1 |s-t|^{\frac{1}{\mu_2}}dtds+\tau_2\int_0^1\int_0^1 \eta^{\frac{s^{\alpha}}{\tau_2}}dtds \\
\leq\ & \mu_1\int_0^1\int_0^1 |s-t|^{\frac{1}{\mu_1}}dtds+\mu_2\int_0^1\int_0^1 |s-t|^{\frac{1}{\mu_2}}dtds+\tau_1\int_0^1\int_0^1 \eta^{\frac{\alpha t}{\tau_1}}dtds+\tau_2\int_0^1\int_0^1 \eta^{\frac{\alpha s}{\tau_2}}dtds \\
=\ & \frac{2\mu_1^3}{(2\mu_1+1)(\mu_1+1)}+\frac{2\mu_2^3}{(2\mu_2+1)(\mu_2+1)}+\tau_1\frac{\eta\left(\frac{\alpha}{\tau_1},\frac{\alpha}{\tau_1}\right)-1}{\ln\eta\left(\frac{\alpha}{\tau_1},\frac{\alpha}{\tau_1}\right)}+\tau_2\frac{\eta\left(\frac{\alpha}{\tau_2},\frac{\alpha}{\tau_2}\right)-1}{\ln\eta\left(\frac{\alpha}{\tau_2},\frac{\alpha}{\tau_2}\right)}
\end{aligned} \tag{2.12}$$

from (2.9)-(2.12), which completes the proof. □

Corollary 7. *Under the assumptions of Theorem 4, and* $\mu=\mu_1=\mu_2>0$, $\tau=\tau_1=\tau_2>0$ *with* $\mu+\tau=1$, *then we have*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\leq\begin{cases}\frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left[\frac{4\mu^3}{(2\mu+1)(\mu+1)}+2\tau\right], & \eta=1\\ \frac{(b-a)}{2}\left|f'\left(\frac{b}{m}\right)\right|^m\left[\frac{4\mu^3}{(2\mu+1)(\mu+1)}+2\tau\frac{\eta\left(\frac{\alpha}{\tau},\frac{\alpha}{\tau}\right)-1}{\ln\eta\left(\frac{\alpha}{\tau},\frac{\alpha}{\tau}\right)}\right], & \eta<1\end{cases}$$

References

[1] R.-F. Bai, F. Qi and B.-Y. Xi, Hermite-Hadamard type inequalities for the m- and (α,m)-logarithmically convex functions. *Filomat,* **27** (2013), 1-7.

[2] M.Z. Sarıkaya, E. Set, M.E. Özdemir: New inequalities of Hermite-Hadamard Type, Volume 12, Issue 4, 2009, Art.11, RGMIA Online: http://rgmia.org/papers/v12n4/set2.pdf

[3] J. Hadamard: Étude sur les propriétés des fonctions entières et en particulier d'une fonction considerée par Riemann, J. Math Pures Appl., 58, (1893) 171–215.

[4] D. S. Mitrinović, J. Pečarić and A. M. Fink: Classical and new inequalities in analysis, KluwerAcademic, Dordrecht, 1993.

[5] S. S. Dragomir and C. E. M. Pearce: Selected topics on Hermite-Hadamard inequalities and applications, RGMIA monographs, Victoria University, 2000. [Online: http://www.staff.vu.edu.au/RGMIA/monographs/hermite-hadamard.html].

[6] S.S Dragomir and R.P.Agarwal,Two inequalities for differantiable mappings and applications to special means of real numbers and trapezoidal formula, Appl. Math. Lett., 11(5) (1998), 91-95.

[7] J. E. Pečarić, F. Proschan and Y. L. Tong: Convex Functions, Partial Orderings, and Statistical Applications, Academic Press Inc., 1992.

[8] M. Tunç, E. Yüksel and İ. Karabayır, On some inequalities for functions whose second derivetives absolute values are α-, m-, (α,m)-logarithmically convex, submitted.

□DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, KILIS 7 ARALIK UNIVERSITY, KILIS, 79000, TURKEY.
E-mail address: □`mevluttunc@kilis.edu.tr`

▽THE INSTITUTE FOR GRADUATE STUDIES IN SCIENCES AND ENGINEERING, KILIS 7 ARALIK UNIVERSITY, KILIS, 79000, TURKEY.
E-mail address: ▽`yuksel.ebru90@hotmail.com`